

Flu Treatment and Prevention: What Pharmacists Need to Know

Top Takeaways

- *Start antivirals ASAP for high-risk patients, even if it's after 48 hours.*
- *Stick with oseltamivir for most high-risk patients. Baloxavir is a one-dose option for eligible outpatients.*
- *Save prophylaxis for select exposed patients the flu vaccine isn't protecting...and start it within 48 hours.*

Patients will look to you to sort out if they need a flu antiviral...with an early surge in influenza in some parts of the US.

Treat high-risk patients ASAP. Recommend an antiviral for suspected or confirmed flu in patients at high risk of complications...such as kids under age 5, adults 65 and up, patients who are immunocompromised or pregnant, and those with chronic conditions (diabetes, lung disease, etc).

Don't wait for a positive test when flu is circulating...and don't let a negative rapid antigen test rule out flu. Plus, ensure these patients still get treatment if symptoms started MORE than 48 hours ago.

But treatment is optional for otherwise healthy patients. It can shorten symptoms by about a day if started within 48 hours.

Stick with oseltamivir for most high-risk patients. It has the most data and is preferred in pregnancy, severe illness, and young kids. It costs about \$20 for 5 days. Advise taking it with food...to limit nausea.

Verify adolescent and adult doses...usually 75 mg twice daily for 5 days. Keep in mind to reduce the dose if eGFR is 60 mL/min or less.

Double-check weight-based doses for kids. Don't be surprised to see 3.5 mg/kg twice daily for infants 9 to 11 months. The American Academy of Pediatrics recommends a higher dose than CDC's 3 mg/kg twice daily.

Dispense oseltamivir suspension with an oral syringe...and give a 3 or 5 mL oral syringe for infants instead of the one in the box.

Stay tuned. There's a preprint study that shows oseltamivir isn't effective and may worsen outcomes in patients who are critically ill with influenza...versus placebo. We're awaiting peer review and publication.

Consider baloxavir as a one-dose option. It seems to work as well as oseltamivir for uncomplicated flu...with less nausea. It's approved for patients age 5 and up.

Keep in mind, it's not recommended in pregnancy, breastfeeding, or patients who are severely ill or immunocompromised. Data are lacking.

Tell patients not to take it with dairy, antacids, or supplements with calcium, iron, magnesium, or zinc...these can lower absorption.

Don't expect to see generic baloxavir yet. It's FDA-approved, but launch depends on patent issues. Brand Xofluza costs about \$170 per dose.

Reserve inhaled zanamivir (Relenza) for patients age 7 and up when oral options aren't feasible...and avoid it in asthma or COPD.

Save post-exposure prophylaxis for select patients. Advise it for exposed residents during institutional outbreaks...unvaccinated household contacts of a high-risk person...and patients who are immunocompromised.

Cite this document as follows: Article, Flu Treatment and Prevention: What Pharmacists Need to Know, Pharmacist's Letter, November 2026

The content of this article is provided for educational and informational purposes only, and is not a substitute for the advice, opinion or diagnosis of a trained medical professional. If your organization is interested in an enterprise subscription, email sales@trchealthcare.com.

© 2026 Therapeutic Research Center (TRC). TRC and Pharmacist's Letter and the associated logo(s) are trademarks of Therapeutic Research Center. All Rights Reserved.

Explain that either oseltamivir or baloxavir is effective, but oseltamivir has a longer track record.

Encourage flu vaccines as long as flu is circulating...along with fluids, rest, handwashing, staying home when sick, etc.

Get our chart, *Antivirals for Influenza*, for med considerations and dosing...and use our *Flu Vaccines for 2026-2027* chart for guidance.

Key References:

- Committee on Infectious Diseases. Recommendations for Prevention and Control of Influenza in Children, 2026-2027: Policy Statement. Pediatrics. 2026 Aug 10. doi:10.1542/peds.2026-078778. Epub ahead of print.
- Committee on Infectious Diseases. Recommendations for Prevention and Control of Influenza in Children, 2026-2027: Technical Report. Pediatrics. 2026 Aug 10. doi:10.1542/peds.2026-078779. Epub ahead of print.
- CDC. Influenza antiviral medications: summary for clinicians. March 10, 2026.
<https://www.cdc.gov/flu/hcp/antivirals/summary-clinicians.html> (Accessed October 6, 2026).
- Uyeki TM, Bernstein HH, Bradley JS, et al. Clinical Practice Guidelines by the Infectious Diseases Society of America: 2018 Update on Diagnosis, Treatment, Chemoprophylaxis, and Institutional Outbreak Management of Seasonal Influenza. Clin Infect Dis. 2019 Mar 5;68(6):e1-e47.
- Medication pricing by Elsevier, accessed October 2026.

Pharmacist's Letter. November 2026, No. 421101

Cite this document as follows: Article, Flu Treatment and Prevention: What Pharmacists Need to Know, Pharmacist's Letter, November 2026

The content of this article is provided for educational and informational purposes only, and is not a substitute for the advice, opinion or diagnosis of a trained medical professional. If your organization is interested in an enterprise subscription, email sales@trchealthcare.com.

© 2026 Therapeutic Research Center (TRC). TRC and Pharmacist's Letter and the associated logo(s) are trademarks of Therapeutic Research Center. All Rights Reserved.